

NECESSARY AND SUFFICIENT CONDITIONS FOR MCSHANE'S IDENTITY AND VARIATIONS

SER PEOW TAN, YAN LOI WONG, AND YING ZHANG

ABSTRACT. Greg McShane introduced a remarkable identity for lengths of simple closed geodesics on the once punctured torus with a complete, finite volume hyperbolic structure. Bowditch later generalized this and gave sufficient conditions for the identity to hold for general type-preserving representations of a free group on two generators Γ to $\mathrm{SL}(2, \mathbf{C})$, this was further generalized by the authors to obtain sufficient conditions for a generalized McShane's identity for arbitrary (not necessarily type-preserving) non-reducible representations. In this note we extend the above by giving necessary and sufficient conditions for the generalized McShane identity to hold (Akiyoshi, Miyachi and Sakuma had proved it for type-preserving representations). We also give a version of Bowditch's variation of McShane's identity to once-punctured torus bundles, in the case where the monodromy is generated by a reducible element, and an asymptotic version of this using the generalized Markoff maps interpretation, and provide necessary and sufficient conditions for the variations to hold.

1. Introduction

For a once punctured torus $\mathbb{T}$ with a complete, finite volume hyperbolic structure, McShane showed [8] that the following identity holds:

$$\sum_{\gamma} \frac{1}{1 + e^{l(\gamma)}} = \frac{1}{2}, \quad (1)$$

where γ ranges over all simple closed geodesics on $\mathbb{T}$, and $l(\gamma)$ is the hyperbolic length of γ in $\mathbb{T}$. This identity is independent of the hyperbolic structure on the torus, that is, it holds for all points in the Teichmüller space $\mathcal{T}_{1,1}$ of the punctured torus. It can also be stated in terms of representations of the fundamental group Γ of $\mathbb{T}$ into $\mathrm{SL}(2, \mathbf{R})$.

On the other hand, Bowditch gave an alternative proof of (1) in [3] via Markoff triples, and extended it in [5] to type-preserving representations $\Gamma \rightarrow \mathrm{SL}(2, \mathbf{C})$ which satisfy certain conditions which we call here the BQ-conditions (Definition 2.7). Akiyoshi, Miyachi and Sakuma [1] generalized this (see Proposition 5.2 there, stated here as Theorem 1.3) to give (implicitly) necessary and sufficient conditions for the identity to hold. In this note, we generalize the above to get necessary and sufficient conditions for the generalized McShane identity to hold for arbitrary (non-elementary) representations $\Gamma \rightarrow \mathrm{SL}(2, \mathbf{C})$ (Theorem 1.6). Bowditch also gave variations of McShane's identity for representations corresponding to complete hyperbolic structures on once-punctured torus bundles M over the circle, where the

The authors are partially supported by the National University of Singapore academic research grant R-146-000-056-112. The third author is also partially supported by the National Key Basic Research Fund (China) G1999075104.

monodromy of M is generated by a hyperbolic (Anosov) element of the mapping class group of the once-punctured torus—here the mapping class group is identified with $\mathrm{SL}(2, \mathbf{Z})$. This was generalized in [15] to general representations. We give further variations in the case where the monodromy of M is generated by a reducible element (corresponding to a parabolic element of $\mathrm{SL}(2, \mathbf{Z})$), and where some version of the BQ-conditions holds (Theorem 1.9). Finally, we also give an asymptotic averaging version of this for certain generalized Markoff maps (Theorem 2.6).

Let $\mathcal{X} = \mathrm{Hom}(\Gamma, \mathrm{SL}(2, \mathbf{C}))/\mathrm{SL}(2, \mathbf{C})$ where the quotient is by the conjugation action. To simplify notation, we denote by ρ instead of $[\rho]$ the conjugacy class of a representation; there should be no confusion as we are mostly interested in functions of the trace which are conjugacy invariants. We denote by $\mathcal{X}_{\mathrm{tp}} \subset \mathcal{X}$ the subset of type preserving representations; by definition, these are the representations satisfying $\mathrm{tr}([\rho(a), \rho(b)]) = -2$ for any generating pair a, b of Γ . We define an equivalence relation $\sim$ on Γ by $g \sim h$ if g is conjugate to h or h^{-1} . Then $\Gamma/\sim$ can be identified with the set of free homotopy classes of unoriented closed curves on $\mathbb{T}$. Note that for $\rho \in \mathcal{X}$, the trace function $\mathrm{tr}(\rho[g])$ is well-defined on the classes $[g] \in \Gamma/\sim$. We denote by $\hat{\Omega}$ the subset of $\Gamma/\sim$ which is identified with the set of free homotopy classes of non-trivial, non-peripheral unoriented simple closed curves on $\mathbb{T}$.

Conventions of Notation. Throughout this paper, we always assume that, for $u \in \mathbf{C}$,

- (a) $\sqrt{u}$ has positive real part if $u \notin (-\infty, 0]$ and has positive imaginary part if $u \in (-\infty, 0)$;
- (b) $\log u$ ($u \neq 0$) has imaginary part in $(-\pi, \pi]$; and
- (c) $\cosh^{-1}(u)$ has real part ≥ 0 and imaginary part $\in (-\pi, \pi]$, and has imaginary part $\in [0, \pi]$ when real part = 0.

The letters μ and τ are always frozen for two complex numbers so that $\mu = \tau + 2$. Implicitly, τ is for the trace of the commutator $[A, B] := ABA^{-1}B^{-1}$ of $A, B \in \mathrm{SL}(2, \mathbf{C})$ and μ is for $\mathrm{tr}^2 A + \mathrm{tr}^2 B + \mathrm{tr}^2 AB - \mathrm{tr} A \mathrm{tr} B \mathrm{tr} AB$, and the relation $\mu = \tau + 2$ follows from the well known Fricke trace identity. Consequently, the letter ν is always frozen as $\nu := \cosh^{-1}(-\tau/2) = \cosh^{-1}(1 - \mu/2)$.

Definition 1.1. Define the function $h : \mathbf{C} \setminus \{0\} \rightarrow \mathbf{C}$ by

$$h(x) = \frac{1}{2} \left(1 - \sqrt{1 - \frac{4}{x^2}} \right). \quad (2)$$

Then Bowditch's extension and reformulation of (1), stated in terms of representations, is as follows.

Theorem 1.2. (Theorem 3 [5]) *Suppose $\rho \in \mathcal{X}_{\mathrm{tp}}$ satisfies*

- (i) $\mathrm{tr} \rho[g] \notin [-2, 2]$ for all $[g] \in \hat{\Omega}$; and
- (ii) $|\mathrm{tr} \rho[g]| \leq 2$ for only finitely many classes $[g] \in \hat{\Omega}$.

Then

$$\sum_{[g] \in \hat{\Omega}} h(\mathrm{tr} \rho[g]) = \frac{1}{2}, \quad (3)$$

where the sum converges absolutely.

We call conditions (i) and (ii) in Theorem 1.2 the BQ-conditions. Note that the summands in (3) are equal to the summands in (1) for representations coming from complete finite area hyperbolic structures on $\mathbb{T}$. Bowditch has conjectured that the representations satisfying the BQ-conditions are precisely the quasifuchsian representations (Conjecture A in [5]). Note also that because $h(x)$ is an even function, the theorem holds for representations into $\mathrm{PSL}(2, \mathbb{C})$ as well (the trace of the commutator is well-defined).

Condition (i) implies that there are no elliptics or accidental parabolics in $\rho(\hat{\Omega})$ and condition (ii) is clearly necessary for the sum in (3) to converge. It turns out these conditions are almost necessary; it is easy to see that it is necessary that there are no elliptics (otherwise there are infinitely many $[g] \in \hat{\Omega}$ with $|\mathrm{tr}(\rho[g])| < K$ for some $K > 0$ and the sum diverges), but it turns out that one can allow for (a finite number of) accidental parabolics and (3) will still hold. We have the following result of Akiyoshi, Miyachi and Sakuma [1].

Theorem 1.3. ([1], Proposition 5.2) *Suppose $\rho \in \mathcal{X}_{\mathrm{tp}}$. Then*

$$\sum_{[g] \in \hat{\Omega}} h(\mathrm{tr} \rho[g]) = \frac{1}{2},$$

and the sum converges absolutely, if and only if

- (i') $\mathrm{tr} \rho[g] \notin (-2, 2)$ for all $[g] \in \hat{\Omega}$, and
- (ii) $|\mathrm{tr} \rho[g]| \leq 2$ for only finitely many classes $[g] \in \hat{\Omega}$.

We will see later (Example 4.1) that conjecturally, the extra representations included in Theorem 1.3 come from the points in the Maskit embedding of $\mathcal{T}_{1,1}$ (see [7] for background and a discussion of the geometry of this embedding), and the cusp points on the boundary of the embedding. In particular, there are one or two accidental parabolics in these extremal cases.

We turn now to more general representations. A representation $\rho \in \mathcal{X}$ is said to be a τ -**representation**, where $\tau \in \mathbb{C}$, if for some (hence every) pair of free generators $a, b \in \Gamma$, $\mathrm{tr} \rho([a, b]) = \tau$. Denote by $\mathcal{X}_\tau \subset \mathcal{X}$ the set of τ -representations (so $\mathcal{X}_{\mathrm{tp}} = \mathcal{X}_{-2}$). Note that this makes sense for representations into $\mathrm{PSL}(2, \mathbb{C})$ as well since the trace of the commutator is well-defined. It is well known that $\rho(\Gamma)$ is elementary if and only if $\tau = 2$, which corresponds to reducible representations, and a different type of geometry; this will be explored in a future paper. Hence, for now, we will only be interested in τ -representations with $\tau \neq 2$. To state the generalized version of McShane's identity formulated in terms of representations, we need the following definition.

Definition 1.4. For a fixed $\tau \neq \pm 2$, let $\nu = \cosh^{-1}(-\tau/2)$. Define a function $\mathfrak{h}_\tau : \mathbb{C} \setminus \{\pm\sqrt{\tau+2}\} \rightarrow \mathbb{C}$ by

$$\mathfrak{h}_\tau(x) = \log \frac{1 + (e^\nu - 1)h(x)}{1 + (e^{-\nu} - 1)h(x)}, \quad (4)$$

where $h(x)$ is given as in (2).

Remark 1.5. It can be verified by an elementary calculation that

$$\mathfrak{h}_\tau(\mathrm{tr} \rho[g]) = 2 \tanh^{-1} \left(\frac{\sinh \nu}{\cosh \nu + e^{l(\rho[g])}} \right),$$

where $l(\rho[g])$ is the complex translation length of $\rho[g]$. This was the version used in [13], and has a more geometric flavor.

Theorem 1.6. *Let $\rho : \Gamma \rightarrow \mathrm{SL}(2, \mathbf{C})$ be a τ -representation, where $\tau \neq \pm 2$, and let $\nu = \cosh^{-1}(-\tau/2)$. Then*

$$\sum_{[g] \in \hat{\Omega}} \mathfrak{h}_\tau(\mathrm{tr} \rho[g]) = \nu \pmod{2\pi i}, \quad (5)$$

and the sum converges absolutely, if and only if

- (i') $\mathrm{tr} \rho[g] \notin (-2, 2)$ for all $[g] \in \hat{\Omega}$, and
- (ii) $|\mathrm{tr} \rho[g]| \leq 2$ for only finitely many classes $[g] \in \hat{\Omega}$.

Remark 1.7. Note that $\mathfrak{h}_\tau$ is not defined at $\pm\sqrt{\tau+2}$, but this does not cause any essential restriction since it was shown in [15] that if $\mathrm{tr} \rho[g] = \pm\sqrt{\tau+2}$ for some $[g] \in \hat{\Omega}$, then part(ii) of the BQ-conditions is not satisfied. It was also shown in [15] that the BQ-conditions are sufficient for (5) to hold with absolute convergence; Theorem 1.6 is an extension of the result there. To get a feeling for the theorem, the reader should consider representations arising from hyperbolic structures on the three-holed sphere (pair of pants) with geodesic boundary—they satisfy the BQ-conditions—and then consider deformations where some, or all of the boundary components degenerate to cusps; see Example 4.2 (these satisfy conditions (i') and (ii) in Theorem 1.6). See also [6] for a more detailed analysis of all such real representations.

To state our next result, we consider the action of the mapping class group of $\mathbb{T}$, $\mathcal{MCG} \cong \mathrm{SL}(2, \mathbf{Z})$, and its induced action on Γ and $\mathcal{X}_\tau$. Recall that $\mathcal{MCG}$ is the group of isotopy classes of diffeomorphisms of $\mathbb{T}$ which fixes the puncture (that is, each element of $\mathcal{MCG}$ fixes a neighborhood of the puncture pointwise). Every $H \in \mathcal{MCG}$ induces an automorphism H_* of Γ and H acts on $\mathcal{X}_\tau$ by

$$H(\rho)(g) = \rho(H_*(g))$$

for $\rho \in \mathcal{X}_\tau$ and $g \in \Gamma$. Bowditch [4] studied representations $\rho \in \mathcal{X}_{\mathrm{tp}}$ stabilized by a cyclic subgroup $\langle H \rangle < \mathcal{MCG} \cong \mathrm{SL}(2, \mathbf{Z})$ generated by a hyperbolic element and proved a variation of the McShane's identity. This was subsequently generalized in [15]. The result (again restated here in terms of representations) is as follows.

Theorem 1.8. (Theorem A, [4]; see also [15]) *Suppose that $\rho \in \mathcal{X}_\tau$ ($\tau \neq 2$) is stabilized by $\langle H \rangle < \mathcal{MCG} \cong \mathrm{SL}(2, \mathbf{Z})$, where H is a hyperbolic element of $\mathrm{SL}(2, \mathbf{Z})$. Suppose*

- (i) $\mathrm{tr} \rho[[g]] \notin [-2, 2]$ for all $[[g]] \in \hat{\Omega}/\langle H_* \rangle$, and
- (ii) $|\mathrm{tr} \rho[[g]]| \leq 2$ for only finitely many classes $[[g]] \in \hat{\Omega}/\langle H_* \rangle$.

Then

$$\sum_{[[g]] \in \hat{\Omega}/\langle H_* \rangle} \mathfrak{h}(\mathrm{tr} \rho[[g]]) = 0 \pmod{2\pi i}, \quad (6)$$

where if $\tau = -2$ then $\mathfrak{h}(x) = h(x)$, and if $\tau \neq -2$ then $\mathfrak{h}(x) = \mathfrak{h}_\tau(x)$. Moreover, the sum converges absolutely.

Using essentially the same arguments, we can relax the condition (i) to:

- (i') $\mathrm{tr} \rho[[g]] \notin (-2, 2)$ for all $[[g]] \in \hat{\Omega}/\langle H_* \rangle$,

as in Theorems 1.3 and 1.6. However, it is not difficult to see that for fixed τ , the set of $\rho \in \mathcal{X}_\tau$ stabilized by $\langle H \rangle$ is finite (so there is a rigidity-type property), so it is possible that this does not give any new examples of representations satisfying this variation of the identity. Instead, we prove the following version of the above where the stabilizer $\langle H \rangle$ is generated by a reducible (parabolic) element H , and the deformation space is more interesting. In this case the induced action H_* of H on Γ is given by $H_*(a) = a^{\pm 1}$, $H_*(b) = a^N b^{\pm 1}$, $N \in \mathbf{Z}$, for some fixed generating pair a, b of Γ .

Theorem 1.9. *Suppose that $\rho \in \mathcal{X}_\tau$ ($\tau \neq 2$) is stabilized by $\langle H \rangle < \mathcal{MCG} \cong \mathrm{SL}(2, \mathbf{Z})$, where H is a parabolic element of $\mathrm{SL}(2, \mathbf{Z})$. Let $[g_0]$ be the unique element in $\hat{\Omega}$ fixed by H_* . Then*

$$\sum_{[[g]] \in \hat{\Omega}/\langle H_* \rangle - [[g_0]]} \mathfrak{h}(\mathrm{tr}\rho[[g]]) = 0 \pmod{2\pi i}, \quad (7)$$

where $\mathfrak{h} = h$ if $\tau = -2$, and $\mathfrak{h} = \mathfrak{h}_\tau$ if $\tau \neq -2$, and the sum converges absolutely, if and only if

- (i) $\mathrm{tr}\rho[[g]] \notin (-2, 2)$ for all $[[g]] \in \hat{\Omega}/\langle H_* \rangle - [[g_0]]$, and
- (ii) $|\mathrm{tr}\rho[[g]]| \leq 2$ for only finitely many classes $[[g]] \in \hat{\Omega}/\langle H_* \rangle$.

The above can also be stated in terms of once punctured torus bundles over the circle with monodromy a reducible element of $\mathcal{MCG}$. In this case $\hat{\Omega}/\langle H_* \rangle$ can be identified with the set of free homotopy classes of non-trivial, non-peripheral simple closed curves on the fibre, and $\hat{\Omega}/\langle H_* \rangle - [[g_0]]$ with the subset which excludes the class of the fixed curve. Note that there is no generalization of Theorem B of [4] in this case, because $\hat{\Omega}/\langle H_* \rangle - [[g_0]]$ is essentially one-sided. It is also not difficult to see that, for $\mathcal{X}_{\mathrm{tp}}$, the remaining case of representations stabilized by a subgroup generated by an elliptic element H gives rise to either representations which satisfy the BQ-conditions, or the representation corresponding to the trivial Markoff map, that is, where $\mathrm{tr}\rho(g) = 0$ for all $g \in \Gamma$. Hence Theorems 1.8 and 1.9 together with the observation above give a rather complete answer to the cases of representations in $\mathcal{X}_{\mathrm{tp}}$ stabilized by either finite or (virtually) cyclic subgroups. In a future paper [16] we will explore representations in $\mathcal{X}_{\mathrm{tp}}$ stabilized by subgroups which are not finite or virtually cyclic, and also the end invariants of a generalized Markoff map ϕ . It turns out that the extended Bowditch's conditions are more useful there. We defer the statement of our final result (Theorem 2.6) to the next section as it is more conveniently stated in terms of generalized Markoff maps.

We give a brief discussion of the proofs. Following Bowditch [5] [4], the results above are reformulated (see §2) in terms of Markoff maps (for Theorem 1.3); generalized Markoff maps (for Theorem 1.6); and generalized Markoff maps stabilized by a parabolic element of $\mathrm{PSL}(2, \mathbf{Z})$ (for Theorem 1.9). The basic idea in [5], extended in [15], was that the BQ-conditions imply that the (generalized) Markoff maps have Fibonacci growth either on Ω (for Theorems 1.3 and 1.6) or on suitable branches of the tree Σ (for Theorem 1.9), which was sufficient to give the absolute convergence of the sums in Theorems 1.3, 1.6 and 1.9. The actual sum was then computed by a tricky computation, using the ψ -functions $\frac{x}{yz}$, $\frac{y}{zx}$ and $\frac{z}{xy}$ associated to a Markoff triple (x, y, z) , or the functions $\Psi(y, z, x)$, $\Psi(z, x, y)$, $\Psi(x, y, z)$ associated to a generalized Markoff triple (x, y, z) . The existence of accidental parabolics (regions where

the generalized Markoff map takes values ± 2) means that the associated generalized Markoff map does not have Fibonacci growth. However, it turns out that the condition of Fibonacci growth is stronger than is necessary for the absolute convergence of the sums and the computation of their values. By analyzing the behavior of a generalized Markoff map on the regions surrounding a region with value ± 2 , we show that the absolute convergence of the sums with the resulting values in Theorems 1.3, 1.6 and 1.9 still holds in the presence of a finite number of accidental parabolics, so that the BQ-conditions can be relaxed (see §3, and also [1] where the proof follows along very similar lines). There are some subtleties involved in the proof of Theorem 1.9 for some special cases; these are taken care of in Lemma 3.13. The proof of Theorem 2.6 is similar to that for Theorem 1.9. Because of the non-periodicity, an asymptotic type identity is probably the best we can hope for in this case. Finally, to illustrate some of the extremal cases covered by our results, we give some examples in §4.



Remark 1.10. McShane's original identity (1) has also been generalized to more general hyperbolic surfaces with cusps by McShane himself [9], to hyperbolic surfaces with cusps and/or geodesic boundary components by Mirzakhani [11], to hyperbolic surfaces with cusps, geodesic boundary and/or conical singularities, and to classical Schottky groups by the authors in [13], [14]. It would be interesting to extend the results here to representations corresponding to these generalizations.

Acknowledgements. We would like to thank Makoto Sakuma for his encouragement and for bringing our attention to Proposition 5.2 of [1]. We would like to apologize to the authors of [1] for the oversight in a previous version of this paper.

2. Generalized Markoff Maps

In this section we reformulate Theorems 1.3, 1.6 and 1.9 in terms of (generalized) Markoff maps, following [5], [4] and [15], and also state our last result, Theorem 2.6. The discussion will be brief. The reader is referred to the references listed above for details.

Let Σ be a binary tree properly embedded in the (hyperbolic) plane. A *complementary region* of Σ is the closure of a connected component of the complement. Denote by $V(\Sigma)$, $E(\Sigma)$ and $\Omega(\Sigma)$ the set of vertices, edges, and complementary regions of Σ respectively, or simply, just V , E and Ω respectively. We will fix a concrete realization of the above concepts by thinking of Σ as the dual to the Farey tessellation $\mathcal{F}$ of the hyperbolic plane (upper half plane) by ideal triangles. Recall that $\frac{p}{q}, \frac{p'}{q'} \in \mathbf{Q} \cup \{\infty\}$ (here $\infty := \frac{1}{0}$) with $\gcd(p, q) = \gcd(p', q') = 1$ are Farey neighbors if $|pq' - p'q| = 1$, and that the Farey triangulation consists of edges which are complete hyperbolic geodesics joining all pairs $\{\frac{p}{q}, \frac{p'}{q'}\} \subset \mathbf{Q} \cup \{\infty\}$ which are Farey neighbors, see Figure 1. In this way, there is a natural action of $\mathrm{PSL}(2, \mathbf{Z})$ on Σ , also on Ω , and there is a natural correspondence of Ω with $\mathbf{Q} \cup \{\infty\}$, which embeds as a dense subset of $\mathbf{R} \cup \{\infty\} \cong S^1$, and inherits the cyclic ordering from S^1 (S^1 can be identified with the projective lamination space of $\mathbb{T}$). Indeed, with this identification, the action of $\mathrm{PSL}(2, \mathbf{Z})$ on $\Omega \leftrightarrow \mathbf{Q} \cup \{\infty\}$ is the usual one. We use the letters $X, Y, Z, W, \dots$ to denote the elements of Ω , and also introduce the notation $X_{\frac{p}{q}}$ to indicate that $X_{\frac{p}{q}} \in \Omega$ corresponds to $\frac{p}{q} \in \mathbf{Q} \cup \{\infty\}$, where we use


 FIGURE 1. Farey tessellation and the binary tree Σ .

 FIGURE 2. The directed edge $\vec{e} = (X, Y; Z \rightarrow W)$

$\frac{1}{0}$ to denote ∞ , and $\frac{p}{q}$ (with $p \in \mathbf{Z}$, $q \in \mathbf{N}$, $\gcd(p, q) = 1$) to denote elements of $\mathbf{Q}$. We use the notation $e \leftrightarrow (X, Y; Z, W)$ to indicate that $e = X \cap Y$ and $e \cap Z$ and $e \cap W$ are the endpoints of e . Denote by $\vec{E}(\Sigma)$ (or just $\vec{E}$) the set of directed edges of Σ where the direction is always taken to be from the tail to the head (as in the direction of the arrow). We also use e to denote the underlying undirected edge corresponding to the directed edge $\vec{e}$, and $-\vec{e}$ to denote the directed edge in the opposite direction of $\vec{e}$. And for a directed edge $\vec{e} \in \vec{E}(\Sigma)$, we use $\vec{e} = (X, Y; Z \rightarrow W)$ to indicate that $e \cap W$ is the head of $\vec{e}$, that is, $\vec{e}$ is the directed edge from Z to W , as shown in Figure 2.

Definition 2.1. For a complex number $\mu \in \mathbf{C}$, a μ -Markoff triple is an ordered triple $(x, y, z) \in \mathbf{C}^3$ of complex numbers satisfying the μ -Markoff equation:

$$x^2 + y^2 + z^2 - xyz = \mu. \quad (8)$$

Thus Markoff triples are just 0-Markoff triples. It is easily verified that if (x, y, z) is a μ -Markoff triple, so are $(x, y, xy - z)$, $(x, xz - y, z)$, $(yz - x, y, z)$ and the permutation triples of each of them.

Definition 2.2. A map $\phi : \Omega \rightarrow \mathbf{C}$ is a μ -Markoff map if

- (i) for every vertex $v \in V(\Sigma)$, the triple $(\phi(X), \phi(Y), \phi(Z))$ is a μ -Markoff triple, where $X, Y, Z \in \Omega$ are the three regions meeting v ; and
- (ii) for every edge $e \in E(\Sigma)$ such that $e \leftrightarrow (X, Y; Z, W)$, we have

$$xy = z + w, \quad (9)$$

where $x = \phi(X), y = \phi(Y), z = \phi(Z), w = \phi(W)$.

As in [15] we shall use Φ_μ to denote the set of all μ -Markoff maps, so that the set of all Markoff maps is denoted by Φ_0 in our notation. We shall also use the following convention of Bowditch [5].

Convention. We always use upper case letters denote the regions and the corresponding lower case letters denote the values of ϕ on the regions, that is, $\phi(X) = x, \phi(Y) = y, \phi(Z) = z$, etc.

As in the case of Markoff maps, if the edge relation (9) is satisfied along all edges, then it suffices that the vertex relation (8) be satisfied at a single vertex. In fact one may establish a bijective correspondence between μ -Markoff maps and μ -Markoff triples, by fixing three regions X, Y, Z which meet at some vertex v ; for concreteness, let us say $X = X_{\frac{0}{1}}, Y = X_{\frac{1}{1}}$ and $Z = X_{\frac{1}{0}}$. The correspondence is then given by $\iota : \phi \mapsto (\phi(X_{\frac{0}{1}}), \phi(X_{\frac{1}{1}}), \phi(X_{\frac{1}{0}}))$. This process may be inverted to obtain ι^{-1} by constructing a tree of μ -Markoff triples as Bowditch did in [5] for Markoff triples: Given a μ -Markoff triple (x, y, z) , we set $\phi(X_{\frac{0}{1}}) = x, \phi(X_{\frac{1}{1}}) = y, \phi(X_{\frac{1}{0}}) = z$, and extend over Ω as dictated by the edge relation (9). In this way one obtains an identification of Φ_μ with the variety in $\mathbf{C}^3$ given by the μ -Markoff equation. In particular, Φ_μ gets a nice topology as a subset of $\mathbf{C}^3$.

The natural action of $\mathrm{PSL}(2, \mathbf{Z})$ on Σ and on Ω induces an action on Φ_μ , where if $H \in \mathrm{PSL}(2, \mathbf{Z})$ then $H : \Phi_\mu \rightarrow \Phi_\mu$ is given by

$$H(\phi)(X) = \phi(H(X)), \quad X \in \Omega.$$

There is also an action of the Klein-four group, $\mathbf{Z}_2^2$, on Φ_μ which reverses the signs of two of the values in each triple, and relates representations into $\mathrm{SL}(2, \mathbf{C})$ with representations into $\mathrm{PSL}(2, \mathbf{C})$; however, this will not concern us here.

The key point to obtain the reformulation is that there is also a natural correspondence between conjugacy classes of $(\mu - 2)$ -representations $\rho \in \mathcal{X}_{\mu-2}$ and μ -Markoff maps, once the four regions X, Y, Z, W corresponding to a fixed edge $e \leftrightarrow (X, Y; Z, W)$ are identified with generators a, b, ab and ab^{-1} for Γ respectively. This works because the relations (8) and (9) are equivalent to the trace relations. This also gives a natural identification of Ω with $\hat{\Omega}$ (recall that $\hat{\Omega} \subset \Gamma / \sim$ is the subset corresponding to the non-trivial, non-peripheral simple closed curves on $\mathbb{T}$) and the action of the mapping class group $\mathcal{MCG} \cong \mathrm{SL}(2, \mathbf{Z})$ on $\mathcal{X}_{\mu-2}$ descends to the $\mathrm{PSL}(2, \mathbf{Z})$ action on Φ_μ described earlier. Theorems 1.3, 1.6 and 1.9 can then be reformulated in terms of μ -Markoff maps as follows (we use the same definitions for the functions h and $\mathfrak{h}_\tau$ as in the original statements).

Theorem 2.3. (Reformulation of Theorem 1.3) *Suppose $\phi \in \Phi_0$. Then*

$$\sum_{X \in \Omega} h(\phi(X)) = \frac{1}{2}, \quad (10)$$

and the sum converges absolutely, if and only if

- (i') $\phi(X) \notin (-2, 2)$ for all $X \in \Omega$, and
- (ii) $|\phi(X)| \leq 2$ for only finitely many $X \in \Omega$.

Theorem 2.4. (Reformulation of Theorem 1.6) Suppose $\phi \in \Phi_{\tau+2}$ ($\tau \neq \pm 2$) and let $\nu = \cosh^{-1}(-\tau/2)$. Then

$$\sum_{X \in \Omega} \mathfrak{h}_\tau(\phi(X)) = \nu \pmod{2\pi i}, \quad (11)$$

and the sum converges absolutely, if and only if

- (i') $\phi(X) \notin (-2, 2)$ for all $X \in \Omega$, and
- (ii) $|\phi(X)| \leq 2$ for only finitely many $X \in \Omega$.

Theorem 2.5. (Reformulation of Theorem 1.9) Suppose that $\phi \in \Phi_{\tau+2}$ ($\tau \neq 2$) is stabilized by $\langle H \rangle < \text{PSL}(2, \mathbf{Z})$, where H is a parabolic element of $\text{PSL}(2, \mathbf{Z})$. Let X_0 be the unique element in Ω fixed by H . Then

$$\sum_{[X] \in \Omega/\langle H \rangle - [X_0]} \mathfrak{h}(\phi[X]) = 0 \pmod{2\pi i}, \quad (12)$$

(where $\mathfrak{h} = h$ if $\tau = -2$ and $\mathfrak{h} = \mathfrak{h}_\tau$ if $\tau \neq -2$) and the sum converges absolutely, if and only if

- (i) $\phi[X] \notin (-2, 2)$ for all $[X] \in \Omega/\langle H \rangle - [X_0]$, and
- (ii) $|\phi[X]| \leq 2$ for only finitely many classes $[X] \in \Omega/\langle H \rangle$.

Note that in Theorem 2.5 above, we have $\phi(X_0) = 2 \cos(\frac{2}{p}\pi)$ for some $\frac{2}{p} \in \mathbf{Q}$. See Lemma 3.11 for a proof. Now suppose that $\phi(X_0) \in (-2, 2)$ and $\phi(X_0) \neq 2 \cos(\frac{2}{p}\pi)$ for any $\frac{2}{p} \in \mathbf{Q}$. Suppose further that for each directed edge $\vec{e}$ meeting X_0 only at its head, ϕ satisfies the extended BQ-conditions on $\Omega^{0-}(\vec{e})$ (see Definition 2.7 below). Then we have, in a special case, the following asymptotic averaging and ergodic version of Theorem 2.5.

Theorem 2.6. Suppose $\phi \in \Phi_\mu$ ($\mu \neq 4$) is such that, under a specific identification of Ω with $\mathbf{Q} \cap \{\frac{1}{0}\}$, $\Omega_\phi(2) = \{X_{\frac{1}{0}}\}$ and $\phi(X_{\frac{1}{0}}) \in (-2, 2)$. Then

$$\lim_{N \rightarrow \infty} \Re \left(\frac{\sum_{r \in \mathbf{Q} \cap (0, N]} \mathfrak{h}(\phi(X_r))}{N} \right) = 0, \quad (13)$$

and there exists an increasing sequence $(N_k)_{k=1}^\infty$ of positive integers such that

$$\lim_{k \rightarrow \infty} \sum_{r \in \mathbf{Q} \cap (0, N_k]} \mathfrak{h}(\phi(X_r)) = 0 \pmod{2\pi i}, \quad (14)$$

where $\mathfrak{h} = h$ if $\mu = 0$, and $\mathfrak{h} = \mathfrak{h}_{\mu-2}$ if $\mu \neq 0$.


We end this section by giving the formal definition and notation for generalized Markoff maps satisfying the BQ-conditions or the extended BQ-conditions.

Definition 2.7. First, we denote by $(\Phi_\mu)_Q$ the subset of Φ_μ satisfying the BQ-conditions, that is, $\phi \in (\Phi_\mu)_Q$ if

- (i) $\phi(X) \notin [-2, 2]$ for all $X \in \Omega$, and
- (ii) $|\phi(X)| \leq 2$ for only finitely many $X \in \Omega$.

Next, we denote by $(\Phi_\mu)_{\overline{Q}}$ the subset of Φ_μ satisfying the extended BQ-conditions, that is, $\phi \in (\Phi_\mu)_{\overline{Q}}$ if

- (i') $\phi(X) \notin (-2, 2)$ for all $X \in \Omega$, and

FIGURE 3. Two simple circular sets $C(T)$

(ii) $|\phi(X)| \leq 2$ for only finitely many $X \in \Omega$.

Finally, we say that ϕ satisfies the BQ-conditions (resp. the extended BQ-conditions) on a subset Ω' of Ω if (i) (resp. (i')) and (ii) above hold for all $X \in \Omega'$.

3. Proofs of Theorems

We first give a quick sketch of how the corresponding results were obtained for (generalized) Markoff maps ϕ satisfying the BQ-conditions. We need to recall some definitions from [5] and [15].

Recall that $E = E(\Sigma)$ and $\vec{E} = \vec{E}(\Sigma)$ are the set of edges and directed edges of Σ respectively, where the direction for a directed edge is taken to be from the tail to the head. For $e \leftrightarrow (X, Y; Z, W) \in E$, define $\Omega^0(e) = \{X, Y\}$. For $\vec{e} \leftrightarrow (X, Y; Z \rightarrow W) \in \vec{E}$, $\Sigma \setminus \text{int}(e)$ consists of two components, denoted by $\Sigma^+(\vec{e})$ and $\Sigma^-(\vec{e})$, where $\Sigma^+(\vec{e})$ is the component containing the head of $\vec{e}$ and $\Sigma^-(\vec{e})$ is the component containing the tail of $\vec{e}$ (so that Z meets $\Sigma^-(\vec{e})$ and W meets $\Sigma^+(\vec{e})$). Define $\Omega^+(\vec{e})$ (resp. $\Omega^-(\vec{e})$) to be the set of regions in Ω whose boundaries lie entirely in $\Sigma^+(\vec{e})$ (resp. $\Sigma^-(\vec{e})$). Hence $\Omega = \Omega^+(\vec{e}) \sqcup \Omega^0(e) \sqcup \Omega^-(\vec{e})$. Define $\Omega^{0+}(\vec{e}) = \Omega^0(e) \cup \Omega^+(\vec{e})$ (resp. $\Omega^{0-}(\vec{e}) = \Omega^0(e) \cup \Omega^-(\vec{e})$).

In terms of a concrete identification of Ω with $\mathbf{Q} \cup \{\infty\}$, we have the following interpretation of the above definitions. If $\vec{e} = (X_{r_1}, X_{r_2}; X_{r_3} \rightarrow X_{r_4}) \in \vec{E}$, then $\Omega^0(e) \leftrightarrow \{r_1, r_2\}$ (note that r_1 and r_2 are Farey neighbors). Also, r_1, r_2 divides $\mathbf{R} \cup \{\infty\} \cong S^1$ into two open intervals $I^+(\vec{e})$ and $I^-(\vec{e})$ where $r_3 \in I^-(\vec{e})$ and $r_4 \in I^+(\vec{e})$. (Bowditch [5] calls the closure of $I^-(\vec{e})$ the impression of the branch $\Omega^-(\vec{e})$.) Then $\Omega^-(\vec{e}) \leftrightarrow (\mathbf{Q} \cup \{\infty\}) \cap I^-(\vec{e})$. For example, in Figure 1, if $\vec{e} = (X_{\frac{1}{2}}, X_{\frac{1}{4}}; X_{\frac{2}{3}} \rightarrow X_{\frac{1}{3}})$, then $\Omega^0(\vec{e}) = \{X_{\frac{1}{2}}, X_{\frac{1}{4}}\}$, and $\Omega^-(\vec{e}) = \{X_{\frac{2}{q}} : \frac{1}{2} < \frac{2}{q} < \frac{1}{3}\}$. In other words, to each directed edge $\vec{e}$, there corresponds an open interval $I^-(\vec{e})$ of S^1 with rational endpoints which are Farey neighbors. This can be visualized as the tail of a “comet” (where one thinks of the directed edge as a comet); $\Omega^-(\vec{e})$ is then the “rational dust” in this tail.

For a given subtree $T \subseteq \Sigma$, the set $C(T) \subseteq \vec{E}$ is the subset of directed edges $\vec{e}$ such that $\vec{e} \in C(T)$ if and only if $e \cap T$ consists of only one point, the point being the head of $\vec{e}$. If T is finite, then so is $C(T)$. For example, if T consists of a single vertex v , then $C(T)$ consists of the three directed edges with heads at v , and if T

consists of a single edge and its two vertices, then $C(T)$ consists of four directed edges, as shown in Figure 3. A finite subset C of $\vec{E}$ is said to be a *circular set* if it has the form $C(T)$ for some finite subtree T of Σ . It is not difficult to see, using the concrete realization, that C is either circular or of the form $\{\vec{e}, -\vec{e}\}$ for some $\vec{e} \in \vec{E}$ if and only if

$$\bigcup_{\vec{e} \in C} \overline{I^-(\vec{e})} = \mathbf{R} \cup \{\infty\} \cong S^1, \quad \text{and}$$

$$I^-(\vec{e}_1) \cap I^-(\vec{e}_2) = \emptyset, \quad \text{for } \vec{e}_1, \vec{e}_2 \in C, \vec{e}_1 \neq \vec{e}_2.$$

Using the analogy of comets as directed edges, this corresponds to a finite set of comets whose tails fill the entire horizon at infinity, and such that no comet lies in the “shadow” of another.

Now let us fix one $\phi \in (\Phi_\mu)_Q$. (We shall deal with Theorems 1.8 and 2.5 later.) So ϕ satisfies the BQ-conditions (see Definition 2.7). Using ϕ we can define a map

$$\alpha = \alpha_\phi : E \rightarrow \vec{E}$$

such that, for $e \leftrightarrow (X, Y; Z, W)$,

$$\alpha(e) = (X, Y; Z \rightarrow W)$$

if $|\phi(Z)| > |\phi(W)|$ and one may choose an arbitrary direction if $|\phi(Z)| = |\phi(W)|$. Thus the direction of the edge is always from the region with greater absolute value of ϕ to the region with smaller absolute value of ϕ .

There is a way to measure the growth rate of the function $\log^+ |\phi|$ on Ω (where $\log^+ x := \max\{0, \log x\}$), which will be useful for proving absolute convergence of certain series, by comparing it to the Fibonacci function $F : \Omega \rightarrow \mathbf{N}$, which we can take to be

$$F(X_{\frac{p}{q}}) = |p| + |q|, \quad \text{where } \gcd(p, q) = 1.$$

(Strictly speaking, this is the Fibonacci function associated to the edge $e \leftrightarrow (X_{\frac{1}{0}}, X_{\frac{0}{1}}; X_{\frac{-1}{1}}, X_{\frac{1}{1}})$; but it was shown in [5] that in fact the concept of having Fibonacci bounds is independent of the edge used.) Then $\log^+ |\phi|$ has a lower Fibonacci bound on Ω if there exists a constant $\kappa > 0$ such that $\log^+ |\phi(X)| > \kappa F(X)$ for all but finitely many $X \in \Omega$. Clearly, we can talk about lower Fibonacci bounds for $\log^+ |\phi|$ restricted to subsets $\Omega' \subseteq \Omega$ as well. The key point is that if $\log^+ |\phi|$ has a lower Fibonacci bound on $\Omega^-(\vec{e})$ for some $\vec{e} \in \vec{E}$, then the sum $\sum_{X \in \Omega^-(\vec{e})} \mathfrak{h}_\tau(\phi(X))$ converges absolutely (to a value which can be determined).

We now outline (with some slight modifications) in several steps the basic strategy used in [5] and [15] to prove the (generalized) McShane's identity for a fixed $\phi \in (\Phi_\mu)_Q$. For any $k > 0$, we define

$$\Omega(k) := \Omega_\phi(k) = \{X \in \Omega \mid |\phi(X)| \leq k\}.$$

By BQ(ii), $\Omega(2)$ is finite. Let us first suppose that $\Omega(2)$ is non-empty.

Step 1.

Let $X_0 \in \Omega(2)$ and let $\{\vec{e}_n \mid n \in \mathbf{Z}\}$ be the bi-infinite sequence of directed edges in Σ bounding X_0 , say in clockwise order around X_0 , where $\vec{e}_n$ is directed from $\vec{e}_{n-1}$ to $\vec{e}_{n+1}$, and let Y_n be the region sharing the edge e_n with X_0 (see Figure 4 where we have represented the boundary of X_0 by a straight line so all the edges e_n lie on a straight line). Then by BQ(i), $\phi(X_0) \notin [-2, 2]$, so $|\phi(Y_n)|$ grows exponentially




FIGURE 4.

in $|n|$ for $|n|$ sufficiently large (Lemma 3.3 [5], see also [15]). We can find a finite subarc

$$J(X_0) = \bigcup_{-N < i < N} e_i, \quad \text{where } N \text{ is a positive integer}$$

such that

- (i) if $Y_j \in \Omega(2)$ then $e_j \subseteq J(X_0)$;
- (ii) if $e_k \not\subseteq J(X_0)$ then $\alpha(e_k)$ is directed towards $J(X_0)$, that is, $\alpha(e_k) = \vec{e}_k$ for $k \leq -N$ and $\alpha(e_k) = -\vec{e}_k$ for $k \geq N$; and
- (iii) $\log^+ |\phi|$ has a lower Fibonacci bound on both $\Omega^-(\vec{e}_N)$ and $\Omega^-(\vec{e}_{-N})$.

Step 2.

Now we take the union

$$T = \bigcup_{X_0 \in \Omega(2)} J(X_0).$$

It can be shown that T is always connected (so T is a finite subtree of Σ) and further that $\alpha(e)$ is directed towards T for all $e \not\subseteq T$.

Aside 1:

In the case where $\Omega(2) = \emptyset$, T consists of a single vertex v which is a sink, with $\alpha(e_i)$ directed towards v for all the three edges e_1, e_2, e_3 incident to v . In this case $\log^+ |\phi|$ has a lower Fibonacci bound on every $\vec{e} \in \vec{E}$.

Step 3.

Now we want to show that for all $\vec{e} \in C(T)$, $\log^+ |\phi|$ has a lower Fibonacci bound on $\Omega^-(\vec{e})$. There are two possibilities. First, $|\phi(X)| > 2$ for $X \in \Omega^0(\vec{e})$ and furthermore $\alpha(e) = \vec{e}$, then $\log^+ |\phi|$ has a lower Fibonacci bound on $\Omega^-(\vec{e})$ (see Corollary 3.6 [5], see also [15]). The second possibility is that $|\phi(X)| \leq 2$ for exactly one of the $X \in \Omega^0(\vec{e})$. This can only occur in the situation of part (iii) of Step 1, with $\vec{e} = -\vec{e}_N$ or $\vec{e}_{-N}$, so again $\log^+ |\phi|$ has a lower Fibonacci bound on $\Omega^-(\vec{e})$.

Since $\Omega = \bigcup_{\vec{e} \in C(T)} \Omega^0(\vec{e})$, this gives the absolute convergence of the sums in (10) and (11), as the fact that $\log^+ |\phi|$ has a lower Fibonacci bound implies the absolute convergence of the sums.

Aside 2:

A slight modification is necessary to deal with Theorems 1.8 and 2.5. We construct the tree T as before in Step 2, except that now it is not finite. However, the tree is invariant under the action of $\langle H \rangle$, and by taking the quotient of T by the action of $\langle H \rangle$, we do indeed get a finite graph. Following [4], by considering the sum over

suitable branches, we again get the absolute convergence of the sum in (6) and (12), when $\phi/\langle H \rangle$ satisfies the BQ-conditions.

Step 4.

To get the actual values of the sums requires the introduction of a function

$$\Psi_\mu : \{\mu\text{-Markoff triples}\} \rightarrow \mathbf{C}$$

(where $\mu = \tau + 2 \neq 0, 4$) defined by

$$\Psi_\mu(x, y, z) := \log \frac{1 + (e^\nu - 1)(z/xy)}{\sqrt{1 - \mu/x^2} \sqrt{1 - \mu/y^2}}, \quad (15)$$

where $\nu = \cosh^{-1}(1 - \mu/2)$, for a μ -Markoff triple (x, y, z) with $x, y \neq 0, \pm\sqrt{\mu}$.

Remark 3.1. In the case where $\mu = 0$ if we set $\mu = 0$ (and hence $\nu = 0$) in (15) we would have $\Psi_0(x, y, z) \equiv 0$; in this case we define a function Ψ'_0 by

$$\Psi'_0(x, y, z) = z/xy,$$

which is used extensively by Bowditch [5] and is in fact the partial derivative of $2\Psi_\mu(x, y, z)$ with respect to ν evaluated at $\nu = 0$ (or equivalently, $\mu = 0$).

Although the function Ψ_μ is rather complicated, what concerns us most will be the properties of the function and the way it relates to the sums in (6), (10), (11), and (12).

Lemma 3.2. (Properties of the function Ψ_μ) *Suppose $\mu \neq 0, 4$.*

(i) *If (x, y, z) is a μ -Markoff triple such that $x, y, z \neq 0, \pm\sqrt{\mu}$ then*

$$\Psi_\mu(x, y, z) + \Psi_\mu(y, z, x) + \Psi_\mu(z, x, y) = \nu \pmod{2\pi i}. \quad (16)$$

(ii) *If (x, y, z) and (x, y, w) are μ -Markoff triples such that $x, y \neq 0, \pm\sqrt{\mu}$ and $z + w = xy$ then*

$$\Psi_\mu(x, y, z) + \Psi_\mu(x, y, w) = \nu \pmod{2\pi i}. \quad (17)$$

Definition 3.3. For a fixed $\phi \in \Phi_\mu$ ($\mu \neq 0, 4$), we define the *edge-weight function* ψ associated to ϕ by

$$\psi(\vec{e}) := \psi_\phi(\vec{e}) = \Psi_\mu(x, y, z), \quad (18)$$

where $\vec{e} = (X, Y; W \rightarrow Z)$ (that is, Z is at the head of $\vec{e}$) and, by our convention, $x = \phi(X), y = \phi(Y), z = \phi(Z)$.

Remark 3.4. In the case where $\mu = 0$ we define

$$\psi(\vec{e}) = z/xy \quad (19)$$

as explained before.

Then we have the following properties, proven in [15] with the help of Lemma 3.2 above.

Lemma 3.5. (Properties of the edge-weight function ψ) *Suppose $\phi \in \Phi_\mu$ (where $\mu \neq 0, 4$) and $\phi(X) \neq 0, \pm\sqrt{\mu}$ for all $X \in \Omega$. Then*

(i) *for $\vec{e} \in \vec{E}$,*

$$\psi(\vec{e}) + \psi(-\vec{e}) = \nu \pmod{2\pi i}; \quad (20)$$

(ii) for a circular set $C \subset \vec{E}$,

$$\sum_{\vec{e} \in C} \psi(\vec{e}) = \nu \pmod{2\pi i}. \quad (21)$$

Remark 3.6. In the case where $\mu = 0$, Lemmas 3.2 and 3.5 hold with ν replaced by 1 and with $\text{mod } 2\pi i$ removed, as used by Bowditch in [5].

Definition 3.7. (The function $\hat{\mathfrak{h}}_\tau$) For $\tau \in \mathbf{C} \setminus \{\pm 2\}$, we set $\mu = \tau + 2$ and define another function $\hat{\mathfrak{h}}_\tau : \mathbf{C} \setminus \{\pm \sqrt{\mu}\} \rightarrow \mathbf{C}$, which is a specific half, modulo $2\pi i$, of the function $\mathfrak{h}_\tau$ defined before, by

$$\hat{\mathfrak{h}}_\tau(x) = \log \frac{1 + (e^\nu - 1)h(x)}{\sqrt{1 - \mu/x^2}}, \quad (22)$$

where $\nu = \cosh^{-1}(-\tau/2)$ and h is the function defined by (2).

It can be checked by direct calculation that

$$\mathfrak{h}_\tau = 2\hat{\mathfrak{h}}_\tau \pmod{2\pi i}. \quad (23)$$

We also have

$$\hat{\mathfrak{h}}_\tau(x) = \lim_{y \rightarrow \infty} \Psi_\mu(x, y, z), \quad (24)$$

where $\mu = \tau + 2$ and z is given by

$$z = \frac{xy}{2} \left(1 - \sqrt{1 - 4 \left(\frac{1}{x^2} + \frac{1}{y^2} - \frac{\mu}{x^2 y^2} \right)} \right). \quad (25)$$

Note that the properties (23) and (24) are crucial for the following theorem to hold, as can be seen in [15].

Theorem 3.8. Suppose $\phi \in \Phi_\mu$ ($\mu = \tau + 2 \neq 0, 4$) and $\vec{e} \in \vec{E}$ is such that $\log^+ |\phi|$ has a lower Fibonacci bound on $\Omega^-(\vec{e})$. Then


$$\sum_{X \in \Omega^0(e)} \hat{\mathfrak{h}}_\tau(\phi(X)) + \sum_{X \in \Omega^-(\vec{e})} \mathfrak{h}_\tau(\phi(X)) = \psi(\vec{e}) \pmod{2\pi i}, \quad (26)$$

where the infinite sum converges absolutely. $\square$

Note that the second sum on the left hand side of (26) is an infinite sum while the first sum has only two summands.

Putting everything together, we can now prove Theorems 2.3, 2.4 and 2.5 in the case where the BQ-conditions are satisfied. We just split the infinite sum in each case into a finite number of infinite sums over the tails of a circular set of directed edges as in Step 3, evaluate each using Theorem 3.8, and then apply (ii) of Lemma 3.5 to get the sum. A slight variation is needed for Theorem 2.5: we need to choose a suitable tree T' which is a “fundamental domain” for the invariant tree T and we use the fact that there are two edges $\vec{e}', \vec{e}''$ in the circular set of $C(T')$ with $\psi(\vec{e}'') = \psi(-\vec{e}')$ because H maps $\vec{e}'$ to $-\vec{e}''$ (see [4] for details).

Now a careful examination of the above sketch will show that the proof in the general case breaks down only at Step 1(iii); specifically, only in the last statement there. This is because if we have $\phi(X_0) = \pm 2$ for some $X_0 \in \Omega$, and suppose $Y_n, n \in \mathbf{Z}$ are the neighboring regions of X_0 , then $|\phi(Y_n)|$ only grows linearly in


 FIGURE 5. $\phi(X_0) = 2$

$|n|$ for large $|n|$, and not exponentially as in the case when $\phi(X_0) \notin [-2, 2]$. So for the directed edges $-\vec{e}_N$ and $\vec{e}_{-N}$ directed towards $J(X_0)$ in Step 1(iii), $\log^+ |\phi|$ does not have lower Fibonacci bounds on $\Omega^-(-\vec{e}_N)$ and $\Omega^-(\vec{e}_{-N})$. Nonetheless, we show in the following proposition that the sums in (10), (11) and (12) over $\Omega^{0-}(-\vec{e}_N)$ and $\Omega^{0-}(\vec{e}_{-N})$ converge absolutely, and with appropriate adjustments, to the values $\psi(-\vec{e}_N)$ and $\psi(\vec{e}_{-N})$ as in the case when $\phi(X_0) \notin [-2, 2]$.

Proposition 3.9. *Suppose $\phi \in \Phi_\mu$ ($\mu = \tau + 2 \neq 0, 4$) and $\vec{e} \in \vec{E}$ with $\Omega^0(e) = \{X_0, Y_0\}$ such that $\phi(X_0) = \pm 2$ and $\Omega(2) \cap \Omega^{0-}(\vec{e}) = \{X_0\}$. Then*

$$\sum_{X \in \Omega^0(e)} \hat{\mathfrak{h}}_\tau(\phi(X)) + \sum_{X \in \Omega^-(\vec{e})} \mathfrak{h}_\tau(\phi(X)) = \psi(\vec{e}) \pmod{2\pi i} \quad (27)$$

where the infinite sum converges absolutely.

Proof. For simplicity of notation, let us write $\hat{\mathfrak{h}} = \hat{\mathfrak{h}}_\tau$, $\mathfrak{h} = \mathfrak{h}_\tau$ in this proof.

We may assume $\phi(X_0) = 2$ since the proof for the case where $\phi(X_0) = -2$ is essentially the same. The proof given here is very similar to that given in [1] for the Φ_0 case, although somewhat more complicated because of the nature of $\mathfrak{h}$. Let $Y_n, n \geq 1$ be the sequence of neighboring regions of X_0 which lie in $\Omega^-(\vec{e})$ and let $e_n, n \geq 0$ be the edge $Y_n \cap X_0$ (so $e_0 = e$). Let $\vec{e}_n, n \geq 1$ be the directed edge on e_n which is directed towards e . Let $v_n, n \geq 1$ be the vertex between $\vec{e}_{n-1}$ and $\vec{e}_n$. Let $\varepsilon_n, n \geq 1$ be the edge $Y_{n-1} \cap Y_n$ and $\vec{\varepsilon}_n \in \vec{E}$ be the directed edge on ε_n with its head at v_n (see Figure 5).

Let $y_n = \phi(Y_n), n \geq 0$. Then an easy computation using (8) and (9) gives

$$y_n = y_0 + n(\mu - 4)^{1/2},$$

for some choice of the square root. Since $\mu \neq 4$, y_n grows linearly in n .

Since $\Omega(2) \cap \Omega^{0-}(\vec{\varepsilon}_n) = \emptyset$, we know that $\log^+ |\phi|$ has a lower Fibonacci bound on $\Omega^{0-}(\vec{\varepsilon}_n)$ and hence $\sum_{X \in \Omega^{0-}(\vec{\varepsilon}_n)} \mathfrak{h}(\phi(X))$ converges absolutely for all $n \geq 1$.

We need to show that $\sum_{X \in \Omega^{0-}(\vec{e})} |\mathfrak{h}(\phi(X))|$ converges. This follows from the following lemma since

$$\Omega^{0-}(\vec{e}) = \{X_0\} \cup \bigcup_{n=1}^{\infty} \Omega^{0-}(\vec{\varepsilon}_n).$$

Lemma 3.10. *There exists a constant $\kappa > 0$ such that for sufficiently large n*

$$\sum_{X \in \Omega^{0-}(\vec{\varepsilon}_n)} |\mathfrak{h}(\phi(X))| \leq \kappa \cdot \frac{1}{n^2}. \quad (28)$$

Proof. There is a constant $\kappa_1 > 0$ such that for all $n \geq 1$, $|y_n| \geq \kappa_1 n$ and hence $\log |y_n| \geq \log n + \log \kappa_1$.

Let F_{ε_n} , $n \geq 1$ be the Fibonacci function with respect to edge ε_n (see the beginning of this section where the Fibonacci function was defined for the edge between $X_{\frac{1}{0}}$ and $X_{\frac{0}{1}}$, see also [5]). Hence there exist constants $\kappa_2, \kappa_3, \kappa_4 > 0$ such that if n is sufficiently large, then

$$\log |\phi(X)| \geq \min\{\log |y_{n-1}| - \log 2, \log |y_n| - \log 2\} F_{\varepsilon_n}(X) \quad (29)$$

$$\geq (\log n + \log \kappa_2) F_{\varepsilon_n}(X) \quad (30)$$

$$\geq (\log n + \log \kappa_3) + \kappa_4 F_{\varepsilon_n}(X) \quad (31)$$

for all $X \in \Omega^{0-}(\vec{\varepsilon}_n)$. Thus there exists a constant $\kappa_5 > 0$ so that for sufficiently large n ,

$$\sum_{X \in \Omega^{0-}(\vec{\varepsilon}_n)} |\phi(X)|^{-2} \leq \kappa_5 \cdot \frac{1}{n^2}. \quad (32)$$

This gives (28), noticing that $|\mathfrak{h}(\phi(X))| = O(|\phi(X)|^{-2})$. Actually, since $h(x) = O(x^{-2})$ as $x \rightarrow \infty$, we have

$$\begin{aligned} \mathfrak{h}(x) &= \log \left(\frac{1 + (e^\nu - 1)h(x)}{1 + (e^{-\nu} - 1)h(x)} \right) \\ &= \log \left(1 + \frac{2(\sinh \nu)h(x)}{1 + (e^{-\nu} - 1)h(x)} \right) \\ &\sim 2(\sinh \nu)h(x) \\ &= O(x^{-2}). \end{aligned}$$

This proves Lemma 3.10. $\square$

Thus the infinite sum in (27) converges absolutely. Its actual value can be obtained by sharpening the proof of (26) in [14]. Alternatively, we can also evaluate it using (26) by a simple limit process as follows.

Let $T = \bigcup_{i=1}^{n-1} e_i$. Applying Lemma 3.5(ii) to the circular set $C(T) \subset \vec{E}$ gives

$$\sum_{\vec{e}' \in C(T)} \psi(\vec{e}') = \nu \pmod{2\pi i}.$$

Note that $C(T) = \{-\vec{e}, \vec{\varepsilon}_1, \dots, \vec{\varepsilon}_n, \vec{e}_n\}$. Thus by Lemma 3.5(i) and Theorem 3.8 we have (noticing that $2\hat{\mathfrak{h}} = \mathfrak{h}$)

$$\begin{aligned} \psi(\vec{e}) &= \sum_{i=1}^n \psi(\vec{\varepsilon}_i) + \psi(\vec{e}_n) \pmod{2\pi i} \\ &= \sum_{i=1}^n \left(\hat{\mathfrak{h}}(\phi(Y_{i-1})) + \hat{\mathfrak{h}}(\phi(Y_i)) + \sum_{X \in \Omega^{0-}(\vec{\varepsilon}_i)} \mathfrak{h}(\phi(X)) \right) + \psi(\vec{e}_n) \pmod{2\pi i} \\ &= \hat{\mathfrak{h}}(\phi(Y_0)) + \hat{\mathfrak{h}}(\phi(Y_n)) + \sum_{i=1}^{n-1} \mathfrak{h}(\phi(Y_i)) + \\ &\quad + \sum_{i=1}^n \sum_{X \in \Omega^{0-}(\vec{\varepsilon}_i)} \mathfrak{h}(\phi(X)) + \psi(\vec{e}_n) \pmod{2\pi i}. \end{aligned}$$

Then (27) follows by letting $n \rightarrow \infty$:

$$\begin{aligned}
 \psi(\vec{e}) &= \hat{\mathfrak{h}}(\phi(Y_0)) + \lim_{n \rightarrow \infty} \hat{\mathfrak{h}}(\phi(Y_n)) + \sum_{i=1}^{\infty} \mathfrak{h}(\phi(Y_i)) + \\
 &\quad + \sum_{i=1}^{\infty} \sum_{X \in \Omega^-(\vec{e}_i)} \mathfrak{h}(\phi(X)) + \lim_{n \rightarrow \infty} \psi(\vec{e}_n) \pmod{2\pi i} \\
 &= \hat{\mathfrak{h}}(\phi(Y_0)) + \sum_{X \in \Omega^-(\vec{e})} \mathfrak{h}(\phi(X)) + \hat{\mathfrak{h}}(\phi(X_0)) \pmod{2\pi i} \\
 &= \sum_{X \in \Omega^0(e)} \hat{\mathfrak{h}}(\phi(X)) + \sum_{X \in \Omega^-(\vec{e})} \mathfrak{h}(\phi(X)) \pmod{2\pi i}
 \end{aligned}$$

since it can be easily checked that

$$\lim_{n \rightarrow \infty} \hat{\mathfrak{h}}(\phi(Y_n)) = 0, \quad (33)$$

$$\lim_{n \rightarrow \infty} \psi(\vec{e}_n) = \hat{\mathfrak{h}}(2) = \hat{\mathfrak{h}}(\phi(X_0)). \quad (34)$$

Proposition 3.9 is thus proved. $\square$

Proof of Theorems 2.3 and 2.4. The proofs for Theorems 2.3 and 2.4 now pretty much follow in the same way as the proof for the cases satisfying the BQ-conditions. We construct the finite tree T as in Step 2; the only difference here is that the circular set $C(T) \subset \vec{E}$ may have some directed edges $\vec{e}$ such that $\log^+ |\phi|$ does not have lower Fibonacci bounds on $\Omega^-(\vec{e})$. These edges are adjacent to regions on which ϕ takes the value ± 2 . We then apply Lemma 3.9 to complete the proofs. $\square$

Lemma 3.11. *Suppose a nontrivial μ -Markoff map $\phi \in \Phi_\mu$ is invariant under the action of a parabolic subgroup $\langle H \rangle$ of $\text{PSL}(2, \mathbf{Z})$ where $H \in \text{PSL}(2, \mathbf{Z})$ fixes region $X_0 \in \Omega$ and is conjugated to $\begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$ with $p > 1$. Then $\phi(X_0) = 2 \cos \frac{q\pi}{p}$ for an integer q with $0 \leq q < p$.*

Proof. Let $Y_n, n \in \mathbf{Z}$ be the bi-infinite sequence of neighboring regions of X_0 . Let $x_0 = \phi(X_0) = \lambda + \lambda^{-1}$. If $x_0 \neq \pm 2$, then there exist $A, B \in \mathbf{C}$ such that $AB = (x_0^2 - \mu)/(x_0^2 - 4)$ and $y_n = \phi(Y_n) = A\lambda^n + B\lambda^{-n}$. Now we may suppose H acts on Ω by $H(Y_n) = Y_{n+p}$. Hence we have $y_{-p} = y_0 = y_p$, that is,

$$A\lambda^{-p} + B\lambda^p = A + B = A\lambda^p + B\lambda^{-p}. \quad (35)$$

Since ϕ is nontrivial, A and B are not both 0. Then it is easy to derive from (35) that we always have $\lambda^n = 1$ or $\lambda^n = -1$, and the latter case only occurs when $A + B = 0$. Hence $\phi(X_0)$ has the desired form. $\square$

Remark 3.12. From the above proof we know that if $y_n \neq 0$ for all $n \in \mathbf{Z}$, then we will have $x_0 = \phi(X_0) = 2 \cos \frac{q\pi}{p}$ with $0 \leq q < p$.

Proof of Theorem 2.5. By Lemma 3.11, in this case $\phi(X_0) = 2 \cos(2q\pi/p)$ for some integers $p > 1$ and $0 < q < p$ with $\gcd(p, q) = 1$. If $\phi(X_0) \neq \pm\sqrt{\mu}$, we can follow the proof for the case where ϕ satisfies the BQ-conditions and use Lemma 3.9 to deal with the directed edges $\vec{e}$ in the circular set for which $\log^+ |\phi|$ does not have lower Fibonacci bounds on $\Omega^-(\vec{e})$. The remaining case to take care of is when

$\phi(X_0) = \pm\sqrt{\mu}$ (in this case we cannot fully apply Theorem 3.8 since $\Psi_\mu(x, y, z)$ is not defined when x^2 or y^2 is equal to μ). This is dealt in the following lemma.

Lemma 3.13. *Suppose $(\phi(X_0))^2 = \mu$ in Theorem 2.5. Let $(Y_n), n \in \mathbf{Z}$ be the bi-infinite sequence of neighboring regions of X_0 . Let $\vec{\varepsilon}_n, n \in \mathbf{Z}$ be the directed edge on $\varepsilon_n = Y_{n-1} \cap Y_n$ with its head on X_0 . Then*

$$\sum_{k=1}^{2p} \psi(\vec{\varepsilon}_k) = 0 \pmod{2\pi i}, \quad (36)$$

where $p > 1$ is the index of $\langle H \rangle$ as a subgroup in a maximal parabolic subgroup of $\text{PSL}(2, \mathbf{Z})$, that is, H is conjugated to $\begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$.

Proof. In this case $\phi(X_0) = 2\cos(q\pi/p)$ and $\mu = 4\cos^2(q\pi/p)$. It can be shown that

$$\nu = \cosh^{-1}(1 - \mu/2) = (\pi - 2q\pi/p)i \quad \text{or} \quad (2q\pi/p - \pi)i, \quad (37)$$

depending on whether $q \leq p/2$ or $q \geq p/2$. Without loss of generality, we may assume $q \leq p/2$. Hence $\nu = (\pi - \frac{2q\pi}{p})i$ and $-e^\nu = e^{-i\frac{2q\pi}{p}}$.

Following our convention, let $y_n = \phi(Y_n), n \in \mathbf{Z}$. By (8) we have either

$$\frac{y_j}{y_{j-1}} = e^{i\frac{q\pi}{p}} \quad \text{for all } j \in \mathbf{Z} \quad (38)$$

or

$$\frac{y_j}{y_{j-1}} = e^{-i\frac{q\pi}{p}} \quad \text{for all } j \in \mathbf{Z}. \quad (39)$$

In particular,

$$y_{2p+j} = y_j \quad \text{for all } j \in \mathbf{Z}. \quad (40)$$

Recall that

$$\psi(\vec{\varepsilon}_j) = \log \frac{1 + (e^\nu - 1)(x_0/y_{j-1}y_j)}{\sqrt{1 - \mu/y_{j-1}^2} \sqrt{1 - \mu/y_j^2}}. \quad (41)$$

Hence the desired equality (36) holds if and only if

$$\prod_{j=1}^{2p} \frac{1 + (e^\nu - 1)(x_0/y_{j-1}y_j)}{\sqrt{1 - \mu/y_{j-1}^2} \sqrt{1 - \mu/y_j^2}} = 1; \quad (42)$$

if and only if

$$\prod_{j=1}^{2p} \frac{y_{j-1}y_j + (e^\nu - 1)x_0}{y_j^2 - \mu} = 1; \quad (43)$$

if and only if

$$\frac{y_0y_1 + (e^\nu - 1)x_0}{y_1y_2 + (e^{-\nu} - 1)x_0} \frac{y_2y_3 + (e^\nu - 1)x_0}{y_3y_4 + (e^{-\nu} - 1)x_0} \cdots \frac{y_{2p-2}y_{2p-1} + (e^\nu - 1)x_0}{y_{2p-1}y_{2p} + (e^{-\nu} - 1)x_0} = 1 \quad (44)$$

since it can be easily checked that

$$[y_{j-1}y_j + (e^\nu - 1)x_0][y_{j-1}y_j + (e^{-\nu} - 1)x_0] = (y_{j-1}^2 - \mu)(y_j^2 - \mu). \quad (45)$$

Case 1. Suppose (38) holds. Then $e^{-i\frac{q\pi}{p}}y_j = y_{j-1}$ for all $j \in \mathbf{Z}$. Hence

$$(-e^\nu)y_jy_{j+1} = y_{j-1}y_j \quad \text{for all } j \in \mathbf{Z}, \quad (46)$$

and we have

$$\begin{aligned} \text{LHS of (44)} &= \frac{y_0 y_1 + (e^\nu - 1)x_0}{(-e^\nu)(y_1 y_2 + (e^{-\nu} - 1)x_0)} \frac{y_2 y_3 + (e^\nu - 1)x_0}{(-e^\nu)(y_3 y_4 + (e^{-\nu} - 1)x_0)} \\ &\quad \dots \frac{y_{2p-2} y_{2p-1} + (e^\nu - 1)x_0}{(-e^\nu)(y_{2p-1} y_{2p} + (e^{-\nu} - 1)x_0)} \\ &= 1. \end{aligned}$$

Case 2. Suppose (39) holds. Then $e^{-i\frac{q\pi}{p}} y_{j-1} = y_j$ for all $j \in \mathbf{Z}$. Hence

$$(-e^\nu) y_{j-1} y_j = y_j y_{j+1} \quad \text{for all } j \in \mathbf{Z}, \quad (47)$$

and we have (noticing that $y_{2p+j} = y_j$)

$$\begin{aligned} \text{LHS of (44)} &= \frac{y_0 y_1 + (e^\nu - 1)x_0}{(-e^\nu)[y_1 y_2 + (e^{-\nu} - 1)x_0]} \frac{y_2 y_3 + (e^\nu - 1)x_0}{(-e^\nu)[y_3 y_4 + (e^{-\nu} - 1)x_0]} \\ &\quad \dots \frac{y_{2p-2} y_{2p-1} + (e^\nu - 1)x_0}{(-e^\nu)[y_{2p-1} y_{2p} + (e^{-\nu} - 1)x_0]} \\ &= \frac{y_0 y_1 + (e^\nu - 1)x_0}{y_2 y_3 + (e^\nu - 1)x_0} \frac{y_2 y_3 + (e^\nu - 1)x_0}{y_4 y_5 + (e^\nu - 1)x_0} \frac{y_4 y_5 + (e^\nu - 1)x_0}{y_6 y_7 + (e^\nu - 1)x_0} \\ &\quad \dots \frac{y_{2p-4} y_{2p-3} + (e^\nu - 1)x_0}{y_{2p-2} y_{2p-1} + (e^\nu - 1)x_0} \frac{y_{2p-2} y_{2p-1} + (e^\nu - 1)x_0}{y_{2p} y_{2p+1} + (e^\nu - 1)x_0} \\ &= 1. \end{aligned}$$

This proves Lemma 3.13 and hence Theorem 2.5. $\square$

Proof of Theorem 2.6. Let us write X_∞ for $X_{\frac{1}{0}}$, and X_n for $X_{\frac{n}{1}}$, $n \in \mathbf{Z}$. (The reader is referred to Figure 1 for an identification of Ω with $\mathbf{Q} \cap \{\frac{1}{0}\}$.) By our convention, we write $x_\infty = \phi(X_\infty)$ and $x_n = \phi(X_n)$ for $n \in \mathbf{Z}$. Since $\Omega_\phi(2) = \{X_\infty\}$, we know that $|x_n| > 2$ for all $n \in \mathbf{Z}$.

First, we note that $x_n, n \in \mathbf{Z}$ are bounded. Actually, let $x_\infty = \lambda + \lambda^{-1}$; then $|\lambda| = 1$ (since $x_\infty \in (-2, 2)$), and there exist $A, B \in \mathbf{C}$ such that $AB = (x_\infty^2 - \mu)/(x_\infty^2 - 4)$ and $x_n = A\lambda^n + B\lambda^{-n}$ for all $n \in \mathbf{Z}$. Hence $|x_n| \leq |A| + |B|$ for all $n \in \mathbf{Z}$.

Next, we show that $x_n^2, n \in \mathbf{Z}$ are bounded away from μ . Actually, since

$$x_{n-1}^2 + x_n^2 + x_\infty^2 - x_{n-1}x_nx_\infty = \mu \quad \text{and} \quad x_{n-1} + x_{n+1} = x_nx_\infty,$$

we have

$$x_n^2 - \mu = x_{n-1}(x_nx_\infty - x_{n-1}) - x_\infty^2 = x_{n-1}x_{n+1} - x_\infty^2,$$

hence

$$|x_n^2 - \mu| \geq |x_{n-1}| |x_{n+1}| - x_\infty^2 > 4 - x_\infty^2 > 0.$$

Case I. Suppose $x_\infty^2 \neq \mu$.

Let $e_n, n \in \mathbf{Z}$ be the edge $X_\infty \cap X_n$, and let $\vec{e}_n$ be the directed edge on e_n , directed from e_{n-1} to e_{n+1} . Then by (18)

$$\psi(\vec{e}_n) = \log \frac{x_\infty x_n + (e^\nu - 1)x_{n+1}}{\sqrt{x_\infty^2 - \mu} \sqrt{x_n^2 - \mu}}.$$

Now we claim that $\psi(\vec{e}_n), n \in \mathbf{Z}$ are bounded. Actually, it is clear that both $x_\infty x_n + (e^\nu - 1)x_{n+1}$ and $\sqrt{x_\infty^2 - \mu} \sqrt{x_n^2 - \mu}$ are bounded; furthermore, the latter

is bounded away from 0 as just shown. That $x_\infty x_n + (e^\nu - 1)x_{n+1}$ is bounded away from 0 follows from the following identity:

$$[x_\infty x_n + (e^\nu - 1)x_{n+1}][x_\infty x_n + (e^{-\nu} - 1)x_{n+1}] = (x_\infty^2 - \mu)(x_n^2 - \mu). \quad (48)$$

This proves the claim.

We further claim that $\hat{\mathfrak{h}}(x_n), n \in \mathbf{Z}$ are also bounded. Recall that by definition

$$\hat{\mathfrak{h}}(x_n) = \log \frac{1 + (e^\nu - 1)h(x_n)}{\sqrt{1 - \mu/x_n^2}}.$$

It is clear that $1 + (e^\nu - 1)h(x_n), n \in \mathbf{Z}$ are bounded. That they are bounded away from 0 follows from the following identity:

$$[1 + (e^\nu - 1)h(x_n)][1 + (e^{-\nu} - 1)h(x_n)] = 1 - \mu/x_n^2, \quad (49)$$

together with the fact shown earlier that $x_n^2, n \in \mathbf{Z}$ are bounded away from μ . This proves the claim.

Let $\varepsilon_n, n \in \mathbf{Z}$ be the edge $X_{n-1} \cap X_n$ and $\vec{\varepsilon}_n, n \in \mathbf{Z}$ be the directed edge $(X_{n-1}, X_n; X_{\frac{2n-1}{2}} \rightarrow X_\infty)$ on ε_n .

By Proposition 3.8 we have for each $n \in \mathbf{Z}$,

$$\psi(\vec{\varepsilon}_n) = \hat{\mathfrak{h}}(x_{n-1}) + \hat{\mathfrak{h}}(x_n) + \sum_{r \in \mathbf{Q} \cap (n-1, n)} \mathfrak{h}(x_r) \pmod{2\pi i}, \quad (50)$$

and hence (noticing that $2\hat{\mathfrak{h}} = \mathfrak{h} \pmod{2\pi i}$)

$$\sum_{n=1}^N \psi(\vec{\varepsilon}_n) = \hat{\mathfrak{h}}(x_0) + \hat{\mathfrak{h}}(x_N) + \sum_{r \in \mathbf{Q} \cap (0, N)} \mathfrak{h}(x_r) \pmod{2\pi i}. \quad (51)$$

Note that for $N \geq 1$, $C_N = \{\vec{e}_0, \vec{e}_1, \dots, \vec{e}_N, -\vec{e}_N\} \subset \vec{E}(\Sigma)$ is a circular set. Actually, $C_N = C(T_N)$ for the subtree $T_N = \bigcup_{k=1}^{N-1} e_k$ of Σ (when $N = 1$, T is the vertex $v = X_0 \cap X_1 \cap X_\infty$). Hence by (21) and (20), we have

$$\sum_{n=1}^N \psi(\vec{\varepsilon}_n) = \nu - \psi(-\vec{e}_N) - \psi(\vec{e}_0) = \psi(\vec{e}_N) - \psi(\vec{e}_0) \pmod{2\pi i}. \quad (52)$$

It follows from (51) and (52) that

$$\sum_{r \in \mathbf{Q} \cap (0, N]} \mathfrak{h}(x_r) = \hat{\mathfrak{h}}(x_N) - \hat{\mathfrak{h}}(x_0) + \psi(\vec{e}_N) - \psi(\vec{e}_0) \pmod{2\pi i}. \quad (53)$$

Since it is shown that $\psi(\vec{e}_N)$ and $\hat{\mathfrak{h}}(x_N)$, $N \geq 1$ are all bounded, we have

$$\lim_{N \rightarrow \infty} \Re \left(\frac{\sum_{r \in \mathbf{Q} \cap (0, N]} \mathfrak{h}(x_r)}{N} \right) = 0. \quad (54)$$

This proves (13) for the case where $x_\infty^2 \neq \mu$.

Case II. Suppose $x_\infty^2 = \mu$.

In this case $\psi(\vec{e}_n) (= \infty)$ is not defined. We have to evaluate $\sum_{r \in \mathbf{Q} \cap (0, N]} \mathfrak{h}(x_r) \pmod{2\pi i}$ directly. We shall assume the notation used in Case I.

Note that ν is purely imaginary: $\nu = \cosh^{-1}(1 - \mu/2) \in (0, \pi)i$. Since $AB = 0$, we may assume that $x_n = A\lambda^n$ without loss of generality (recall that $x_\infty = \lambda + \lambda^{-1}$). It is easy to check that we have either $-e^\nu = \lambda^{-2}$ or $-e^\nu = \lambda^2$; and hence correspondingly, either (55) or (56) below holds:

$$x_{n-1}x_n + (e^\nu - 1)x_\infty = (-e^\nu)[x_n x_{n+1} + (e^{-\nu} - 1)x_\infty] \quad \text{for all } n \in \mathbf{Z}, \quad (55)$$

$$x_n x_{n+1} + (e^\nu - 1)x_\infty = (-e^\nu)[x_{n-1}x_n + (e^{-\nu} - 1)x_\infty] \quad \text{for all } n \in \mathbf{Z}. \quad (56)$$

When N is even, we have, with similar calculations as in the proof of Lemma 3.13, that either (when $-e^\nu = \lambda^{-2}$ hence (55) holds)

$$\begin{aligned}
 \sum_{n=1}^N \psi(\vec{\varepsilon}_n) &= \log \frac{\sqrt{1 - \mu/x_N^2}}{\sqrt{1 - \mu/x_0^2}} + \log \frac{\prod_{n=1}^N [x_{n-1}x_n + (e^\nu - 1)x_\infty]}{\prod_{n=1}^N (x_n^2 - \mu)} \pmod{2\pi i} \\
 &= \log \frac{\sqrt{1 - \mu/x_N^2}}{\sqrt{1 - \mu/x_0^2}} + \log \left(\frac{x_0x_1 + (e^\nu - 1)x_\infty}{x_1x_2 + (e^{-\nu} - 1)x_\infty} \frac{x_2x_3 + (e^\nu - 1)x_\infty}{x_3x_4 + (e^{-\nu} - 1)x_\infty} \right. \\
 &\quad \left. \dots \frac{x_{N-2}x_{N-1} + (e^\nu - 1)x_\infty}{x_{N-1}x_N + (e^{-\nu} - 1)x_\infty} \right) \pmod{2\pi i} \\
 &= \log \frac{\sqrt{1 - \mu/x_N^2}}{\sqrt{1 - \mu/x_0^2}} + \frac{N}{2}(\nu + \pi i) \pmod{2\pi i}
 \end{aligned} \tag{57}$$

or (when $-e^\nu = \lambda^2$ hence (56) holds)

$$\begin{aligned}
 \sum_{n=1}^N \psi(\vec{\varepsilon}_n) &= \log \frac{\sqrt{1 - \mu/x_N^2}}{\sqrt{1 - \mu/x_0^2}} + \frac{N}{2}(\nu + \pi i) \\
 &\quad + \log \frac{x_0x_1 + (e^\nu - 1)x_\infty}{x_Nx_{N+1} + (e^\nu - 1)x_\infty} \pmod{2\pi i}.
 \end{aligned} \tag{58}$$

When N is odd, we have similar expressions with one more term in each case since

$$\sum_{n=1}^N \psi(\vec{\varepsilon}_n) = \psi(\vec{\varepsilon}_N) + \sum_{n=1}^{N-1} \psi(\vec{\varepsilon}_n)$$

and $N-1$ is even. Hence we still have (54) in this case, noticing that (a) ν is purely imaginary, (b) $x_N^2, N \geq 1$ are bounded away from μ and 0, and (c) $\psi(\vec{\varepsilon}_N), N \geq 1$ are bounded. This proves (13) for the case where $x_\infty^2 = \mu$.

The proof of (14) is similar. Recall that $x_\infty = \lambda + \lambda^{-1} \in (-2, 2)$, hence $|\lambda| = 1$. Thus there exists an increasing sequence $(N_k)_{k \geq 1}$ of positive integers such that $\lim_{k \rightarrow \infty} \lambda^{N_k} = 1$. Therefore we have $\lim_{k \rightarrow \infty} x_{N_k} = x_0$, $\lim_{k \rightarrow \infty} \psi(\vec{\varepsilon}_{N_k}) = \psi(\vec{\varepsilon}_0)$ and $\lim_{k \rightarrow \infty} \hat{\mathfrak{h}}(x_{N_k}) = \hat{\mathfrak{h}}(x_0)$. Hence if $x_\infty^2 \neq \mu$, we have from (53) (with $N = N_k$) that

$$\sum_{r \in \mathbf{Q} \cap (0, N_k]} \mathfrak{h}(x_r) = \hat{\mathfrak{h}}(x_{N_k}) - \hat{\mathfrak{h}}(x_0) + \psi(\vec{\varepsilon}_{N_k}) - \psi(\vec{\varepsilon}_0) \pmod{2\pi i}. \tag{59}$$

It follows that $\lim_{k \rightarrow \infty} \sum_{r \in \mathbf{Q} \cap (0, N_k]} \mathfrak{h}(x_r) = 0 \pmod{2\pi i}$ in the case $x_\infty^2 \neq \mu$.

Now suppose $x_\infty^2 = \mu$. We may assume that $N_k, k \geq 1$ are all even. Then in the case when $-e^\nu = \lambda^{-2}$, we have from (51) and (57) (with $N = N_k$) that

$$\begin{aligned}
 \sum_{r \in \mathbf{Q} \cap (0, N_k]} \mathfrak{h}(x_r) &= \hat{\mathfrak{h}}(x_{N_k}) - \hat{\mathfrak{h}}(x_0) + \log \left[\sqrt{1 - \mu/x_{N_k}^2} / \sqrt{1 - \mu/x_0^2} \right] \\
 &\quad + \log [(-e^\nu)^{N_k/2}] \pmod{2\pi i}.
 \end{aligned} \tag{60}$$

We have again $\lim_{k \rightarrow \infty} \sum_{r \in \mathbf{Q} \cap (0, N_k]} \mathfrak{h}(x_r) = 0 \pmod{2\pi i}$ in this case, noticing that

$$\lim_{k \rightarrow \infty} \log [(-e^\nu)^{N_k/2}] = \lim_{k \rightarrow \infty} [(\lambda^{-2})^{N_k/2}] = \log 1 = 0 \pmod{2\pi i}.$$

The case when $-e^\nu = \lambda^2$ can be proved similarly, this time using (51) and (58). This finishes the proof of (14) and hence also the proof of Theorem 2.6. $\square$

4. Geometric interpretations and examples

In this section we give some examples of (generalized) Markoff maps $\phi \in \Phi_\mu$ which are extremal with respect to the conditions, namely where $\phi(X) = \pm 2$ for a finite number of $X \in \Omega$ (or $[X] \in \Omega/\langle H \rangle$ in the case of Theorem 2.5), and the geometric situations when they occur.

Example 4.1. We start with examples for Theorem 2.3. The Maskit embedding of $\mathcal{T}_{1,1}$ parameterized by the complex number ζ , with corresponding Kleinian group $\Gamma(\zeta)$ as studied by Linda Keen and Caroline Series in [7] corresponds to the 0-Markoff triple $(2, -i\zeta, -i(\zeta + 2))$ and the corresponding 0-Markoff map $\phi_\zeta \in \Phi_0$ (they used the parameter μ instead of ζ ; here we use ζ to avoid confusion as μ is already used for the generalized Markoff maps). These satisfy the conditions of Theorem 2.3 since the number of simple closed geodesics on the hyperbolic 3-manifold $\mathbf{H}^3/\Gamma(\zeta)$ of length $\leq K$, for any $K > 0$ is finite. For example, if we take ζ such that its imaginary part $\Im \zeta \geq 2$, then it is easy to see that the conditions will be satisfied. These examples correspond to points on the boundary of quasifuchsian space $\mathcal{QF}_{1,1}$ for which McShane's identity still holds. More generally, the same is true even when we look at the cusp points on the boundary of the Maskit embedding, these correspond to the case where exactly two curves are pinched, one on each end of the 3-manifold. On the other hand, Minsky's results from [10], see also the references contained therein, imply that for any other boundary point of $\mathcal{QF}_{1,1}$, there is one or two degenerate ends, in which case there are infinitely many homotopy classes of simple closed curves on the torus of bounded length, and hence McShane's identity does not hold. Hence, at least on $\partial(\mathcal{QF}_{1,1})$, one can determine completely the points where McShane's identity holds. Now if Bowditch's conjecture is true, that is, if the set of Markoff maps satisfying the BQ-conditions are exactly those which come from the quasifuchsian representations, this would give a complete geometric description of all representations for which McShane's identity holds, in the type-preserving case.

Example 4.2. We next give an example of an extremal case for Theorem 2.4, in the case where $\mu = 20$. This is the Markoff map $\phi \in \Phi_{20}$ corresponding to the Markoff triple $(-2, -2, -2)$, which arises from the representation coming from the complete, finite volume hyperbolic structure on the thrice-punctured sphere. Note that in this case $|\Omega(2)| = 3$. It seems likely that there is a fixed constant N such that $|\Omega_\phi(2)| \leq N$ for all generalized Markoff maps ϕ for which McShane's identity holds.

Example 4.3. We finally give some examples for Theorem 2.5. Consider the Markoff map $\phi \in \Phi_0$ corresponding to the triple $(0, 2, 2i)$. It is not difficult to verify that this satisfies the conditions of Theorem 2.5 where the stabilizer H of ϕ is conjugate to $\begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix} \in \mathrm{PSL}(2, \mathbf{Z})$. More generally, so does the map $\phi \in \Phi_0$ corresponding to the triple $(0, \zeta, \zeta i)$ for $|\zeta| > 2$. It is not difficult to construct other examples with other stabilizers, by replacing 0 with $2 \cos(\pi/n)$, and the other values appropriately, for example, the μ -Markoff map $\phi \in \Phi_\mu$, where $\mu = 4 \cos^2(\pi/n)$, corresponding to the triple $(2 \cos(\pi/n), \zeta, e^{i\pi/n} \zeta)$ where again $|\zeta| > 2$. It seems

that most of these examples should correspond to representations arising from hyperbolic three orbifolds. However, we are not sure about the detailed geometric interpretations of these generalized Markoff maps.

REFERENCES

- [1] Hirotaka Akiyoshi, Hideki Miyachi and Makoto Sakuma, *A refinement of McShane's identity for quasifuchsian punctured torus groups*, In the Tradition of Ahlfors and Bers, III. The Ahlfors-Bers Colloquium, Oct. 2001, Univ. of Connecticut at Storrs, W. Abikoff, A. Haas (Eds.), Contemporary Mathematics **355**, 21–40, American Mathematical Society, 2004.
- [2] Hirotaka Akiyoshi, Hideki Miyachi and Makoto Sakuma, *Variations of McShane's identity for punctured surface groups*, London Mathematical Society Lecture Notes, Y. Minsky, M. Sakuma & C. Series (Eds.), Cambridge University Press (to appear).
- [3] Brian H. Bowditch, *A proof of McShane's identity via Markoff triples*, Bull. London Math. Soc. **28** (1996), no. 1, 73–78.
- [4] Brian H. Bowditch, *A variation of McShane's identity for once-punctured torus bundles*, Topology **36** (1997), no. 2, 325–334.
- [5] Brian H. Bowditch, *Markoff triples and quasi-Fuchsian groups*, Proc. London Math. Soc. (3) **77** (1998), no. 3, 697–736.
- [6] William M. Goldman, *The modular group action on real $SL(2)$ -characters of a one-holed torus*, Geom. Topol. **7** (2003), 443–486.
- [7] Linda Keen and Caroline Series, *Pleating coordinates for the Maskit embedding of the Teichmüller space of punctured tori*, Topology **32** (1993), no. 4, 719–749.
- [8] Greg McShane, *A remarkable identity for lengths of curves*, Ph.D. Thesis, University of Warwick, 1991.
- [9] Greg McShane, *Simple geodesics and a series constant over Teichmüller space*, Invent. Math. **132** (1998), no. 3, 607–632.
- [10] Yair N. Minsky *The classification of punctured-torus groups*, Ann. of Math. (2) **149** (1999), no. 2, 559–626.
- [11] Maryam Mirzakhani, *Simple geodesics and Weil-Petersson volumes of moduli spaces of bordered Riemann surfaces*, preprint.
- [12] Makoto Sakuma, *Variations of McShane's identity for the Riley slice and 2-bridge links*, Hyperbolic spaces and related topics (Japanese) (Kyoto, 1998), Surikaiseikenkyusho Kokyuroku No. **1104** (1999), 103–108.
- [13] Ser Peow Tan, Yan Loi Wong and Ying Zhang, *Generalizations of McShane's identity to hyperbolic cone-surfaces*, arXiv:math.GT/0404226 v1.
- [14] Ser Peow Tan, Yan Loi Wong and Ying Zhang, *McShane's identity for classical Schottky groups*, arXiv:math.GT/0411628 v1.
- [15] Ser Peow Tan, Yan Loi Wong and Ying Zhang, *Generalized Markoff maps and McShane's identity*, in preparation.
- [16] Ser Peow Tan, Yan Loi Wong and Ying Zhang, *End invariants of generalized Markoff maps and two generator subgroups of $SL(2, \mathbb{C})$* , in preparation.
- [17] Ying Zhang, *Hyperbolic cone-surfaces, generalized Markoff maps, Schottky groups and McShane's identity*, Ph.D. Thesis, National University of Singapore, 2004.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 2 SCIENCE DRIVE 2, SINGAPORE 117543

E-mail address: mattansp@nus.edu.sg; matwyl@nus.edu.sg; scip1101@nus.edu.sg

AND THE THIRD AUTHOR

DEPARTMENT OF MATHEMATICS, YANGZHOU UNIVERSITY, YANGZHOU 225002, P. R. CHINA

E-mail address: yingzhang@yzu.edu.cn